Technical Bulletin

Genes that fit your farm.



AAC Penhold Canada Prairie Spring Red Wheat



Description:

AAC Penhold is a very short strong strawed CPS wheat that has high yield potential, improved protein content and good disease resistance. AAC Penhold is moderately resistant to FHB and has one of the best disease packages in a CPS wheat so it should be well adapted to all growing regions of western Canada especially those areas concerned with lodging and straw management.

Parentage: 5700PR/HY644-BW//HY469

Strengths:

- Grain yield 106% of 5700PR and 97% of 5701PR (average of all sites in 2010 to 2012 High Yield Wheat Registration trials)
- Improved lodging tolerance compared to 5700PR and 5701PR
- Significantly shorter than 5700PR and 5701PR
- Early maturing, 2 days earlier than 5700PR
- Very large seed size and heavy test weight
- Improved protein over 5700PR and 5701PR
- Moderately Resistant (good) to FHB
- Resistant to common bunt and leaf rust
- Moderately resistant to stripe rust and stem rust Ug99 and variants

Neutral Traits:

 Intermediate resistance to leaf spot and loose smut

Weaknesses:

Response to yellow rust has not been consistent

Breeder:

Dr. Richard Cuthbert Swift Current Research and Development Centre Agriculture and Agri-Food Canada Swift Current, SK

Agronomic Tip: AAC Penhold responds favorably to a heavier seeding rate especially in higher moisture areas. For CPS wheat, the optimum seeding rates ranged from 26 to 32 seeds/ft² in the Dark Brown soil zone to 33 to 42 seeds/ft² in the Black soil zone (this can be in the 180lb/ac range.)

The exceptionally strong straw of AAC Penhold can handle higher fertility – follow fertility recommendations for the target yield of your fields.

PBR 91 Protected

2010-2012 High Yield Wheat Cooperative Trials - Registration Data

Variety	Yield (% of Mean of Checks)	Maturity (days)	Lodging 1 = erect 9 = flat	Height (cm)	Test Weight (kg/kl)	Kernel Weight (mg/kernel)	Grain Protein (%)	Falling Number	FHB Resistance Rating
5700 PR	95	107.3	1.5	79.8	77.2	36.8	11.9	357	MS
5701 PR	105	105.7	1.8	81.4	75.6	37.5	12.3	413	MS
AAC Penhold	101	105.3	1.3	72.3	78.4	42.9	12.9	445	MR

2020 Seed Manitoba - CPSR Wheat Comparison

				Maturity	Height		Resistance to:											
	Site Years	Yield	Protein	+/-	+/-	Spike			Loose		Leaf	Stem	Leaf	Stripe				
Variety	Tested	bu/ac	%	99 days	81cm	Awned	Lodging	Sprouting	Smut	Bunt	Spot	Rust	Rust	Rust	FHB			
AC® Carberry	124	67	14.6	2	0	Υ	VG	F	MR	R	MS	MR	R	MR	MR			
AAC Foray VB	31	77	13.1	1	+5	Υ	VG	Р	MS		MS	MR	R	1	I			
AAC Goodwin	33	74	14.1	2	0	Υ	VG	G	MS		I	ı	R	R	I			
AAC Ryley	30	71	13.0	1	+3	Υ	G	G	_	R	MS	R	R	S	MS			
SY Rowyn	34	74	13.5	1	-5	Υ	VG	F	_	S	I	R	R	MR	MR			
AAC Penhold	44	74	13.8	1	-10	Υ	VG	VG		R	I	MR	R	MR	MR			

Lodging Ratings: F=Fair; G=Good; VG=Very Good

Disease Ratings: R=Resistant; MR=Moderately Resistant; I=Intermediate; MS=Moderately Susceptible; S=Susceptible

2020 Varieties of Grain Crops for Saskatchewan – Wheat Comparison

		Yield a	as % of	Carberry					Resista		Relative		Seed	Test					
	Years	Area	Area					Stem	Leaf	Stripe	Loose		Leaf		Maturity	Head	Weight	Weight	Height
Variety	Tested	1 & 2	3 & 4	Irrigation	Protein	Lodging	Sprouting	Rust	Rust	Rust	Smut	Bunt	Spot	FHB	(days)	Awnedness	(mg)	(kg/hl)	(cm)
AC® Carberry	6	100	100		14.5	VG	F	MR	R	MR	MR	R	MS	MR	102	Y	35.8	80.3	83
AAC Foray VB	5	116	120	122	-1.6	F	Р	MR	R	ı	MS	I	MS	_	0	Υ	+8.0	-1.4	+6
AAC Goodwin	4	116	115		-0.7	G	G	ı	R	R	MS			I	-1	Υ	+1.0	+0.2	+2
5700PR	5	107	113	106	-1.3	VG	F	R	I	S	MS	R	MS	MS	-1	Υ	+3.4	-2.3	0
SY Rowyn	4	103	107		-1.1	F	F	R	R	MR		S		MR	-1	Υ	-4.1	-0.5	-5
AAC Penhold	5	108	111	108	-0.8	VG	VG	MR	R	MR	Ī	R		MR	-2	Y	+5.3	-0.2	-9

G=Good; VG=Very Good; F=Fair; P=Poor; VP=Very Poor

2020 Alberta Seed Guide - CWRS Wheat Comparison

	Overal	Overall Yield (1)		Test Yield Category (2)		Maturity					Resis	tance to:	Disease Resistance:					
Variety	All Sites	Station years of testing	Low < 55 bu/ac	Med 55-80 bu/ac	High >80 bu/ac	Rating (Days +/- Carberry)	Protein %	Test Weight (lb/bu)	Kernel Weight g/1000	Height (cm)	Ldg.	Sprout	Loose Smut	Bunt	Stripe Rust	Leaf Spot	FHB	
	Yield as % of AC® Carberry																	
AC® Carberry												ĺ					Ì	
bu/ac	73		41	63	93													
AC® Carberry	100		100	100	100	104	13.9	63	40	79	VG	F	MR	R	MR	MS	MR	
AAC Foray VB	121	41	117	123	123	0	-1.6	63	51	85	G	G	MS	I	MR	MS	ı	
AAC Goodwin	115	48	112	116	117	-1	-0.5	63	41	83	VG	G	MS	MS	R		ı	
AAC Ryley	111	37	108	112	110	-1	-0.5	60	48	82	G	G		R	S	MS	MS	
5700PR	110	117	108	113	109	-1	-1.8	62	42	75	VG	F	MS	R	MS	MS	MS	
SY Rowyn	106	47	102	109	105	-1	-0.9	62	36	77	G	F	-	S	MR		MR	
AAC Penhold	112	63	108	112	114	-2	-1	63	44	73	VG	G		R	MR	ı	MR	

VG = Very Good; G = Good; F = Fair; P = Poor; VP = Very Poor

Disease Ratings: R=Resistant; MR=Moderately Resistant; I=Intermediate; MS=Moderately Susceptible; S=Susceptible